

AMENDMENTIN THE CLAIMS

Please amend the claims according to the following listing of claims and substitute it for all prior versions and listings of claims in the application.

Claims 1-8 (cancelled).

9. (currently amended) A cream for the treatment of dermatitis ~~characterized in that the volume ratios of respective ingredients are~~ comprising in volume percent: main components consisting of ~~[[L]]lightyellow~~ [[S]]sophora ~~[[R]]root~~ (*Sophora flavescens* Ait.), in an amount of 2.7 to 3.3%; ~~[[F]]turmeric~~ (*Curcuma aromatica* Salisb.), in an amount of 1.8 to 2.2%; ~~[[M]]magnolia~~ [[B]]bark (*Magnolia officinalis* Rehd. Et. Wils.), in an amount of 1.8 to 2.2%; ~~[[M]]moutan~~ [[B]]bark (*Paeonia suffruticosa* Andr.), in an amount of 1.8 to 2.2%; ~~[[I]]isatis~~ [[L]]leaf (*Isatis tinctoria* L.), in an amount of 0.9 to 1.1%; and ~~[[B]]borneo~~ camphor ~~[[F]]tree~~ (*Dryobalanops aromatica* Gaertn. f.), in an amount of 0.9 to 1.1%; and auxiliary components comprising: ~~[[B]]baikal~~ [[S]]skullcap (*Scutellaria baicalensis* Georgi.), in an amount of 1.8 to 2.2%; ~~[[A]]amur~~ [[C]]cork ~~[[F]]tree~~ (*Phellodendron amurense* Rupr.), in an amount of 1.8 to 2.2%; ~~[[A]]angelicae~~ [[D]]dahuricae ~~[[R]]root~~ (*Angelica dahurica* Benth. et Hook.), in an amount of 0.9 to 1.1%; ~~[[L]]lemon~~ (*Citrus Limonum*), in an amount of 0 to 3%; ~~[[S]]smartweed~~ (*Polygonum cuspidatum* Sieb. et Zucc.), in an amount of 0 to 1.1%; ~~[[L]]licorice~~ (*Glycyrrhiza uralensis* Fishch.), in an amount of 0 to 0.55%; ~~[[C]]cnidii~~ Rhizoma root (*Ligusticum chuanxiong* Hort.), in an amount of 0.45 to 1.1%; Japanese Angelica ~~[[R]]root~~ (*Angelica*

sinensis (Olive) Diels.) in an amount of 0 to 0.55%; salicylic acid in an amount of 0.45 to 0.55%; resorcinol in an amount of 0.45 to 0.55%; mutton oil in an amount of 2.7 to 3.3%; alcohol in an amount of 2.7 to 3.3%; and white soft paraffin in an amount of 63 to 78%.

10. (currently amended) A cream for the treatment of dermatitis ~~characterized in that the volume ratios of respective ingredients are~~ comprising in volume percent: main components consisting of Lightyellow Ssophora [[R]]root (*Sophora flavescens* Ait.) in an amount of 3%; Turmeric (*Curcuma aromatica* Salisb.) in an amount of 2%; Mmagnolia [[B]]bark (*Magnolia officinalis* Rehd. Et. Wils.) in an amount of 2%; Mmoutan [[B]]bark (*Paeonia suffruticosa* Andr.) in an amount of 2%; [[H]]isatis [[H]]leaf (*Isatis tinctoria* L.) in an amount of 1%; and Bborneo Ccamphor [[T]]tree (*Dryobalanops aromatica* Gaertn. f.) in an amount of 1%; and auxiliary components comprising Bbaikal Sskullcap (*Scutellaria baikcalensis* Georgi.) in an amount of 2%; [[A]]amur [[C]]cork [[T]]tree (*Phellodendron amurense* Rupr.) in an amount of 2%; Aangelicae Ddahuricae [[R]]root (*Angelica dahurica* Benth. Et Hook.) in an amount of 1%; Elemon (*Citrus Limonum*) in an amount of 3%; Ssmartweed (~~p~~*Polygonum cuspidatum* Sieb. Et Zucc.) in an amount of 1%; Hlicorice (*Glycyrrhiza uralensis* Fishch.) in an amount of 0.5%; Ccnidii root Rhizoma (*Ligusticum chuanxiong* Hort) in an amount of 0.5%; Japanese Aangelica [[R]]root (*Angelica sinensis* (Olive) Diels.) in an amount of 0.5%; salicylic acid in an amount of 0.5%; resorcinol in an amount of 0.5%; mutton oil in an amount of 3%; alcohol in an amount of 3%; and white soft paraffin in an amount of 70.5%.

11. (currently amended) A cream for the treatment of dermatitis ~~characterized in that the volume ratios of respective ingredients are~~ comprising in volume percent: main components

consisting of ~~L~~lightyellow ~~S~~sophora ~~[[R]]~~root (*Sophora flavescens* Ait.), in an amount of 2.7 to 3.3%; ~~T~~turmeric (*Curcuma aromatica* Salisb.), in an amount of 1.8 to 2.2%; ~~M~~magnolia ~~[[B]]~~bark (*Magnolia officinalis* Rehd. et. Wils.), in an amount of 1.8 to 2.2%; ~~M~~moutan ~~[[B]]~~bark (*Paeonia suffruticosa* Andr.), in an amount of 1.8 to 2.2%; ~~I~~isatis ~~[[L]]~~leaf (*Isatis tinctoria* L.), in an amount of 0.9 to 1.1%; and ~~B~~borneo ~~C~~camphor ~~[[T]]~~tree (*Dryobalanops aromatica* Gaertn. f.), in an amount of 0.9 to 1.1%; and auxiliary components comprising ~~B~~baikal ~~S~~skullcap (*Scutellaria baikcalensis* Georgi.), in an amount of 1.8 to 2.2%; ~~[[A]]~~amur ~~[[C]]~~cork ~~[[T]]~~tree (*Phellodendron amurense* Rupr.), in an amount of 1.8 to 2.2%; ~~A~~angelicae ~~D~~dahuricae ~~[[R]]~~root (*Angelica dahurica* Benth. et. Hook.), in an amount of 0.9 to 1.1%; ~~L~~lemon (*Citrus Limonum*), in an amount of 0 to 3%; ~~S~~smartweed (*polygonum cuspidatum* Sieb. et Zucc.), in an amount of 0 to 1.1%; ~~L~~licorice (*Glycyrrhiza uralensis* Fishch.), in an amount of 0 to 0.55%; dimethyl sulfoxide, in an amount of 4.5 to 5.5%; salicylic acid, in an amount of 0.45 to 0.55%; resorcinol, in an amount of 0.45 to 0.55%; mutton oil, in an amount of 2.7 to 3.3%; alcohol, in an amount of 2.7 to 3.3%; and white soft paraffin, in an amount of 60 to 73%.

12. (currently amended) A cream for the treatment of dermatitis ~~characterized in that the volume ratios of respective ingredients are~~ comprising, in volume percent: main components consisting of ~~L~~lightyellow ~~S~~sophora ~~[[R]]~~root (*Sophora flavescens* Ait.), in an amount of 3%; ~~T~~turmeric (*Curcuma aromatica* Salisb.), in an amount of 2%; ~~M~~magnolia ~~[[B]]~~bark (~~M~~*magnolia officinalis* Rehd. et. Wils.), in an amount of 2%; ~~M~~moutan ~~[[B]]~~bark (*Paeonia suffruticosa* Andr.), in an amount of 2%; ~~I~~isatis ~~[[L]]~~leaf (*Isatis tinctoria* L.), in an amount of 1%; and ~~B~~borneo ~~C~~camphor ~~[[T]]~~tree (*Dryobalanops aromatica* Gaertn. f.), in an amount of

1%; and auxiliary components comprising ~~B~~baikal ~~S~~skullcap (*Scutellaria baikcalensis* Georgi.)~~;~~in an amount of 2%; ~~A~~amur ~~[[€]]~~cork ~~[[F]]~~tree (*Phellodendron amurense* Rupr.)~~;~~in an amount of 2%; ~~A~~angelicae ~~D~~dahuricae ~~[[R]]~~root (*Angelica dahurica* Benth. et. Hook.)~~;~~in an amount of 1%; ~~L~~emon (*Citrus Limonum*)~~;~~in an amount of 3%; ~~S~~smartweed (~~p~~*Polygonum cuspidatum* Sieb. et Zucc)~~;~~in an amount 1%; ~~L~~icorice (*Glycyrrhiza uralensis* Fishch.)~~;~~in an amount of 0.5%; dimethyl sulfoxide~~;~~in an amount of 5%; salicylic acid~~;~~in an amount of 0.5%; resorcinol~~;~~in an amount of 0.5%; mutton oil~~;~~in an amount of 3%; alcohol~~;~~in an amount of 3%; and white soft paraffin~~;~~in an amount of 66.5%.

13. (cancelled)

14. (cancelled)

15. (currently amended) The cream for the treatment of dermatitis of claim 11, wherein~~the~~
~~Lightyellow Sophora Root (*Sophora flavescens* Ait.), the Turmeric (*Curcuma aromatica* Salisb.), the Magnolia Bark (*Magnolia officinalis* Rehd. Et. Wils.), the Moutan Bark (*Paeonia suffruticosa* Andr.), the Isatis Leaf (*Isatis tinctoria* L.) and the Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn. f.) serve as main components,~~ the dimethyl sulfoxide ~~serves~~
~~as~~ is an accelerating agent for permeation, the salicylic acid and the resorcinol ~~serve as~~ are
 accelerating agents for keratinization, and, wherein the ratios of the main components:
 accelerating agent for skin permeation : accelerating agents for keratinization are 53 to 89% : 8
 to 38% : 6 to 10%.

16. (cancelled)